



(doxycycline, USP) \*30 mg immediate release & 10 mg delayed release beads

(OR-RAY-SHA)

Oracea Capsules are an oral therapy for the treatment of only the inflammatory lesions of rosacea

# Conversation Starters

## Begin the Conversation

Ready to talk about rosacea but not sure what to ask? Bring these questions to your healthcare provider to help get a healthy dialogue started—and don't delay! The best time to seek help is early on when symptoms are still mild.

1. How do ORACEA Capsules work and are they right for me?  
\_\_\_\_\_
2. How do ORACEA Capsules differ from other oral medications used for rosacea?  
\_\_\_\_\_
3. What can I expect from treatment with ORACEA Capsules?  
\_\_\_\_\_
4. What factors—such as food and beverage, environmental or emotional—may be triggering my rosacea?  
\_\_\_\_\_
5. What products do you suggest I use for face wash, moisturizer, sunscreen and makeup to avoid aggravating my rosacea?  
\_\_\_\_\_
6. Add your own questions:  
\_\_\_\_\_



### Important Safety Information

**Indication:** ORACEA® (doxycycline, USP) 40 mg\* Capsules are indicated for the treatment of only inflammatory lesions (papules and pustules) of rosacea in adult patients. ORACEA Capsules do not lessen the facial redness caused by rosacea. **Adverse Events:** In controlled clinical studies, the most commonly reported adverse events (>2%) in patients treated with ORACEA Capsules were nasopharyngitis, sinusitis, diarrhea, hypertension and aspartate aminotransferase increase. **Warnings/Precautions:** ORACEA Capsules should not be used to treat or prevent infections. ORACEA Capsules should not be taken by patients who have a known hypersensitivity to doxycycline or other tetracyclines. ORACEA Capsules should not be taken during pregnancy, by nursing mothers, or during tooth development (up to the age of 8 years). Although photosensitivity was not observed in clinical trials, ORACEA Capsules patients should minimize or avoid exposure to natural or artificial sunlight. The efficacy of ORACEA Capsules treatment beyond 16 weeks and safety beyond 9 months have not been established.

\*30 mg immediate release & 10 mg delayed release beads

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.