

ONCE-DAILY  
**ORACEA®**  
 (doxycycline, USP) 40\* mg Capsules  
\*30 MG IMMEDIATE RELEASE & 10 MG DELAYED RELEASE BEADS  
 [OR-RAY-SHA]

"For once, I made the decision to put myself first, to commit to clearer skin and treat my rosacea through the good and bad days"

**NOW LOOK AT ME**

Not an actual patient



ROSACEA VS ACNE

## COULD THOSE BUMPS AND BLEMISHES BE ROSACEA?

Acne and rosacea are two common skin conditions. This chart will help you identify if those red bumps are rosacea, acne, or a little bit of both

ORACEA® (doxycycline, USP) 40 mg\* Capsules are indicated for the treatment of the bumps and blemishes of rosacea in adult patients. ORACEA Capsules do not lessen the facial redness caused by rosacea

### SYMPTOMS OF ROSACEA



### SYMPTOMS OF ACNE

#### Pimple-like breakouts

But unlike acne, no blackheads occur

#### Eye problems

Including bloodshot eyes, red and swollen eyelids, problems seeing and/or discomfort

#### Facial redness

Near the central part of your face, cheeks, forehead, nose or chin. Redness may come and go or be permanent

#### Visible blood vessels

Caused by broken small blood vessels near the surface of the skin



If you think you have one or more of the above symptoms, book an appointment with your dermatologist today



#### Oily T-zone

Affecting the nose, chin and forehead



#### Uneven skin texture

Bumpy skin texture, due to blemishes and scars



#### Redness

Occurs around breakouts only



#### Presents as pimples

Including blackheads, whiteheads, pimples or deep painful cysts

To support the conversation with your dermatologist, visit:  
[oracea.com/rosacea-discussion-guide-builder](https://oracea.com/rosacea-discussion-guide-builder)

### IMPORTANT SAFETY INFORMATION

**Indication:** ORACEA® (doxycycline, USP) 40 mg\* Capsules are indicated for the treatment of the bumps and blemishes of rosacea in adult patients. ORACEA Capsules do not lessen the facial redness caused by rosacea. **Adverse Events:** In controlled clinical studies, the most commonly reported adverse events (>2%) in patients treated with ORACEA Capsules were nasopharyngitis, sinusitis, diarrhea, hypertension and aspartate aminotransferase increase. **Warnings/Precautions:** ORACEA Capsules should not be used to treat or prevent infections. ORACEA Capsules should not be taken by patients who have a known hypersensitivity to doxycycline or other tetracyclines. ORACEA Capsules should not be taken during pregnancy, by nursing mothers, or during tooth development (up to the age of 8 years). Although photosensitivity was not observed in clinical trials, ORACEA Capsules patients should minimize or avoid exposure to natural or artificial sunlight. The efficacy of ORACEA Capsules treatment beyond 16 weeks and safety beyond 9 months have not been established

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](https://www.fda.gov/medwatch) or call 1-800-FDA-1088

\*30 mg immediate release and 10 mg delayed release beads

### REFERENCES

1. National Rosacea Society. (2013). Acne Or Rosacea? Accurate Diagnosis Is Key To Treatment. Available: <https://www.rosacea.org/rosacea-review/2013/winter/acne-or-rosacea-accurate-diagnosis-is-key-to-treatment>. Last accessed: September 2021. 2. Learnskin. The subtypes of rosacea and how they are different from acne. Available: <https://www.learnskin.com/articles/the-subtypes-of-rosacea-and-how-they-are-different-from-acne>. Last accessed: September 2021.

This is intended for U.S. audiences only. Information in this fact sheet is not intended as medical advice. Talk to your doctor about medical concerns.

**GALDERMA**

EST. 1981

©2021 Galderma Laboratories, L.P. All rights reserved. All trademarks are the property of their respective owners.

Galderma Laboratories, L.P. 14501 N. Freeway, Fort Worth, TX 76177.

US-ORC-2100083